

STUDY PROTOCOL

Open Access



Predicting unplanned hospital readmission in palliative outpatients (PRePP) – study protocol of a longitudinal, prospective study to identify informal caregiver-related and structural predictors

Leopold Hentschel^{1,2*} , André Wellesen³, Luisa Christin Krause¹, Maria von Havranek¹, Michael Kramer⁴, Beate Hornemann², Martin Bornhäuser^{4,5}, Ulrich Schuler¹ and Katharina Schütte¹

Abstract

Background: Although the majority of German patients in a palliative state prefer to die at home, the actual place of death is most often a hospital. Unplanned hospital readmissions (UHA) not only contradict most patients' preferences but also increase the probability of an aggressive end-of-life treatment. As limited knowledge is available which factors contribute to an UHA, the PRePP-project aims to explore predictors related to informal caregivers (IC) as well as medical and structural factors.

Methods: This prospective, observational, mono-centric study will assess structural and medical factors as well as ICs' psychological burden throughout seven study visits. Starting in April 2021 it will consecutively include 240 patients and their respective IC if available. Standardized measures concerning ICs' Quality of Life (WHOQOL-BREF), psychological distress (NCCN-Distress Thermometer), anxiety (GAD-7) and depressiveness (PHQ-9) will be assessed. If participants prefer, assessment via phone, browser-based or paper-based will be conducted. Medical records will provide routinely assessed information concerning patient-related characteristics such as gender, age, duration of hospital stay and medical condition. Nurse-reported data will give information on whether hospitalization and death occurred unexpectedly. Data will be progressed pseudonymized. Multivariable regression models will help to identify predictors of the primary endpoint "unplanned hospital admissions".

Discussion: The PRePP-project is an important prerequisite for a clinical risk assessment of UHAs. Nevertheless, it faces several methodological challenges: as it is a single center study, representativity of results is limited while social desirability might be increased as the study is partly conducted by the treatment team. Furthermore, we anticipated an underrepresentation of highly burdened participants as they might refrain from participation.

Trial registration: This study was retrospectively registered 19 October 2021 at clinicaltrials.gov (NCT05082389). <https://clinicaltrials.gov/ct2/show/NCT05082389>

*Correspondence: Leopold.Hentschel@ukdd.de

¹ University Palliative Center, University Hospital Carl Gustav Carus, TU Dresden, Fetscherstraße 74, 01307 Dresden, Germany

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Keywords: Unplanned hospital admission, Palliative care, Specialized outpatient palliative care, Informal caregiver, Family caregiver, Psychological distress

Background

With evolving treatment options in several diseases, prolonged survival at the cost of a more complex end-of-life treatment has been achieved in the recent past. In Germany, the need of adequate palliative care as well as the number of structures providing palliative care has increased [1]. In 2014, more than 33.000 cases were treated in specialized palliative care units, 30.000 in hospices and almost 87.500 cases were reported by specialized outpatient palliative care teams [2].

Between 52 and 94% of German patients prefer end-of-life care at their home [3, 4] while the actual place of death was home in less than one third of deceased persons [5] in 2009. Specialized outpatient palliative care teams (german abbreviation: *Spezialisierte Ambulante Palliativversorgung*, SAPV) aim to alleviate symptoms and enable incurably ill patients to stay at their familiar surrounding until death by providing sufficient end-of-life care. Evidence suggests various advantages of preventing patients dying in the hospital. Gomes et al. report that home palliative care increases the chance of dying at home and reduces symptom burden in particular for patients with cancer [6]. Quality of life tends to be higher in patients dying at home [7]. In addition, optimal care by informal caregivers (IC) leads to greater satisfaction and less grief on the part of the relatives [8]. Yet there is a limited understanding of comprehensive costs of palliative care [9], caring at home might relieve financial burden off the health care system as inpatient care accounts for more than 70% of costs [10] during a palliative phase.

Unplanned hospital admissions (UHAs) decrease the probability to die at home [11] and were identified as a factor of overly aggressive treatment near the end of life [12]. UHAs occur due to various reasons including physical, psychological and social factors as well as characteristics of professional support [6, 13]. According to Gamblin and colleagues [14], UHAs occur more frequently in patients being released from palliative care unit (compared to medical oncology ward), which is attributed to a more complex burden of symptoms. Evidence derived from lung cancer patients of a single center suggest UHAs are the result of cancer-related symptoms rather than treatment toxicities [15]. Gamblin et al. illustrated that living alone at home or the presence of at least one child at home were associated with death in hospital [14] while proposing that financial precariousness, exhausted or ill caregiver might contribute to readmission. Furthermore, UHAs caused by a medical indication

can also be altered by the psychosocial factors at the patient's place of residence [16]. A post-discharge palliative care consultation was associated with a significantly lower incidence of UHA [14].

The frequency of UHA as well as the place of death varies internationally [17]. These discrepancies can be explained both by different health care systems with a varying extent of hospice and palliative services [18] and by discrepancies of insurance coverage policies for hospital and palliative care [19–21]. The consensus of experts of the EAPC (European Association of Palliative Care) recommends a minimum of one SAPV-Team per 100.000 inhabitants, while in Germany only 0.22 SAPV-team per 100.000 inhabitants is available [22] with huge differences between federal states. In addition to specialized palliative care, there are further structures available in Germany: Alongside general practitioners and specialists, mobile nursing services and voluntary hospice organizations are involved in the care of palliative patients.

However, a major part of the care is provided by informal caregivers. This includes all family members, friends or neighbors who are involved in the care of a person with health-related restrictions usually without receiving payment [23]. In Germany, up to 4.7 million informal caregivers take care of a frail relatives [24], yet there is no structured evidence available how many of them support terminal-ill patients.

Yet recent research suggests that including informal caregivers in the disease management may improve patients' outcomes [25], it might bear several burdens on the informal-caregiver [26]. These include physical symptoms such as decreased immune functioning, fatigue or sleep disturbance as well as high psychological distress, anxiety and depression [27, 28]. It remains unclear too which extent these burdens contribute to UHA as caregivers, often old or frail him-/herself, might be overwhelmed or too exhausted to provide sufficient care at home.

Overall, there is scarce evidence on factors associated with UHAs in palliative outpatients. It seems of utmost importance to prevent UHAs in order to meet patients' preferred place of death, alter support of the professional system according to the informal caregivers needs and provide economic responsible options for the treating hospital. This project will explore the role of medical, structural as well as caregiver-associated factors as predictors for UHA. It is an important prerequisite for both, clinical implications as well as further research activities.

Methods/design

Study design and setting

Starting in April 2021, the PRePP-project is a prospective, observational, longitudinal, single-arm, single center study to identify structural and medical factors as well as those related to a patient's informal caregiver which predict (unplanned) hospital readmission.

Primary study aim is to explore structural, medical as well as IC-reported factors associated with UHAs. Secondary goals include the description of ICs' psychological distress and supportive care needs.

This study is performed at the SAPV-team of a large tertiary cancer center treating about 300–330 palliative patients a year.

Participants and recruitment

The PRePP-project aims to include two samples of participants, patients and their respective care providers. It aims to include consecutively 240 patients and, if present and consenting, their respective primary IC within 18 months.

Eligible patients who are adult (>17 years), suffer from an advanced, life-limiting disease and are treated by the SAPV-team of our institution will be included.

The primary non-professional caregiver is defined as the adult person that provides most time supporting the patient regardless of being biological related or non-related. ICs can participate if they provide informed consent, are mentally and linguistically able to provide answers independently. IC-reported data can only be collected for patients with consenting ICs. Due to this condition the exact number of IC-participants cannot be predicted.

Study visits and data assessment

Data is obtained from three sources: (1) IC-reported, (2) medical-record based and (3) nurse-reported.

Data of consenting ICs will be obtained three times. Medical-record based and nurse reported data will be obtained seven times. Detailed timeline of study visits can be found in Fig. 1.

After informed consent, which will be obtained at the first regular visit of the SAPV-team (T0), the initial visit collects data of all three sources via tablet-PCs provided by SAPV-team at second home visit (T1). Two subsequent study visits after three (T4) and six months (T7) obtain IC-reported data remotely either browser-based, by mail (paper-pencil) or via phone by study staff. The monthly (T1-T7) assessment includes data derived from medical records and nurse-report.

If no ICs are available or refuse to participate, only the patient's medical report, living situation and nurse-reported data will be gathered monthly (T1–T7).

Electronic data is captured applying REDcapp [29, 30].

Ethics and trial registration

Ethical approval is obtained prior to recruitment from local institutional review board (BO-EK-320072020). According to German law (§34SächsKHG), no study specific informed consent must be obtained from patients to use routinely assessed data from medical records. This trial is registered at clinicaltrials.gov (NCT05082389).

Study measures

Informal caregivers willing to participate answer standardized measures on several domains.

Quality of Life will be assessed administering the WHOQOL-BREF [31], a 26-items inventory to assess a person's Quality of Life including the four domains physical health, psychological, social relationships and environment. Higher scores denote higher Quality of Life in this domain.

Psychological distress will be assessed by the NCCN-Distress Thermometer [32] which comprises of a visual analogue scale ranging from zero to ten with higher scores indicating a more severe burden and a checklist of various symptoms. It is extensively used and cut-off data is available for use in an oncological setting [32]. Preliminary, it was developed for use in cancer patients while it might be applicable in other settings as well [33].

Anxiety will be assessed applying the Generalized Anxiety Disorder 7-item scale [34], a seven-items patient-reported questionnaire with sufficient psychometric properties. Scores range from zero to 21 on a single scale with higher scores indicating a higher level of anxiety. Results can be classified in either one of four classes ranging from minimal to severe anxiety symptoms.

Depressiveness will be measured administering the Patient Health Questionnaire [35], which comprises of nine items. A single sum score is calculated which ranges from 0 to 27 with higher scores indicating a higher level of burden. Results are categorized in four classes ranging from no depressive symptoms to severe symptomatology.

Due to the possible influence of the current COVID-19 pandemic, four self-developed questions considering IC's worries regarding an infection, changes in professional support and psychological burden due to preventive measures such as wearing a face mask were administered.

Nurses will answer two surprise questions [36], predicting whether they would be surprised when the patient will die within the next month (first question) and whether an UHA will occur during the next month (second question).

Characteristics of IC such as age, gender, and kind of relation to the patient, duration of daily care, highest education, and employment status will be obtained.




Fig. 1 Timeline of study visits and source of data. Data source red = medical record; blue = family care-giver; green = nursing practitioner

Furthermore, the amount and type of professional support such as mobile nursing teams will be assessed.

Medical records will provide routinely assessed information concerning characteristics of patient such as age, gender, diagnosis, duration of previous hospitalization, additional supportive services, number of visits by SAPV, symptoms status, admission (e.g. reasons, initiating person, admitting physician, planned vs. unplanned, which procedures applied).

Statistical analysis

The primary endpoint "number of unplanned inpatient admissions" is account variable. Univariate Poisson regressions will be used to analyze the association of structural factors and relative-reported factors or structural data with the primary endpoint. With the planned patient number of 240, it is possible to detect changes of 50% (corresponding to rate ratios of 1.5 and greater, given a base rate of 0.85) with a power of 95% if the influence variable is a binary categorical variable with 50% probability of occurrence. For a binary factor with a probability of occurrence of 90%, the detectable rate ratio with 95% power is still 2.0. For continuous, normally distributed influence variables, rate ratios of >1.25 are detectable with 95% power.

Since a large number of factors are to be tested, the adjustment for multiple testing seems reasonable. According to the explorative approach of the study, the control of the false discovery rate is done by Benjamini–Hochberg procedure. *P* values of the univariate regression analyses that are <0.05 after adjustment with the Benjamini–Hochberg procedure are considered significant. In the multivariable model, all parameters whose *p* values from the univariate analyses are <0.1 after adjustment should be analyzed. The *p*-values of the multivariable model will not be adjusted.

Discussion

The PRePP-projects aims to provide an elaborated understanding of factors predicting hospital readmission in palliative outpatients. As scarce knowledge is available on this issue and little comparable studies are available, we face some practical and operational challenges which future studies could keep in mind.

Representativity and generalizability could be limited as the project is a single-arm and single-center study. This may cause selection bias for certain patients' and ICs' characteristics regarding regional and cultural differences. A multicenter study design could reduce this kind of bias.

The different methods of collecting ICs' data could induce varying response behavior. Participants acquired

by mail or browser-based may feel more anonymous causing more reliable answers, compared to phone surveys. Possible bias due to social desirability while communicating directly with the study staff, could occur. Moreover phone surveys performed by different members of the study team could cause interviewer bias. However we decided to use the opportunity of phone surveys to keep response rate higher, considering elderly participants as not as comfortable using online questionnaires and avoid additional effort for them by returning the mail. As a practical issue we anticipate less participation of highly burdened ICs which could lead to an underestimation of participants' burden.

Concerning the questionnaires used, we face a trade-off between diagnostic accuracy on the one hand while reducing the burden for the participants on the other hand. We decided to administer short instruments such as GAD-7 or PHQ-9 as they use a smaller number of questions and are self-administered compared to structured interviews such as the Structured Clinical Interview for DSM-5 [37]. Furthermore, we aim to describe psychological burden rather than diagnose a psychiatric disease which allows for less sophisticated instruments. This is well in line with other groups performing research in palliative care [38].

As this study was conceptualized during the beginning of the COVID-pandemic there were no standardized questions regarding ICs concerns about COVID available. Future research should progress on this issue, e.g. by developing standardized instruments or at least performing psychometric analyses on available questions.

Regardless of aforementioned challenges, the data obtained from the PRePP-Project serves as an important prerequisite for the development of a screening tool to predict readmission which could be administered to patients routinely in clinical practice.

Abbreviations

EAPC: European Association of Palliative Care; GAD-7: Generalized Anxiety Disorder 7-item scale; IC: Informal caregiver; NCCN: National Comprehensive Cancer Network; NP: Nursing practitioner; PHQ-9: Patient Health Questionnaire 9-item scale; SAPV: Specialized outpatient palliative care (German: Spezialisierte Ambulante Palliativversorgung, abbreviation: SAPV); T0-T7: Time; indicates study visit; UHA: Unplanned hospital admission.

Acknowledgements

We are grateful to all patients and their caregivers that decided to support this project. Furthermore, this project received massive support from the staff of our institution's SAPV-team. Without their dedication, this project would not have been realized.

Authors' contributions

LH and KS conceptualized and designed the study. AW implement the electronic data collection tool. AW, LK and MvH monitor data integrity. LK and MvH recruit patients. MK conduct statistics. BH, MB and US provide expertise

in study conduction. LH, KS, LK and MvH wrote the manuscript. All authors read and approved the final manuscript.

Funding

Open Access funding enabled and organized by Projekt DEAL. This project was supported by an unrestricted grant received from the Stiftung Hochschulmedizin Dresden.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author upon reasonable request.

Declarations

Ethics approval and consent to participate

This study was approved by the Ethics Committee of the Technical University Dresden (BO-EK-320072020). The study is conducted in accordance with the Helsinki Declaration. Informed consent was obtained from informal caregiver prior to study inclusion. No study specific informed consent must be obtained from patients as we use routinely assessed data from medical records. Nevertheless, as part of the routine admission process to the SAPV-care, a regarding consent is signed. This trial is registered at clinicaltrials.gov (NCT05082389).

Consent for publication

Not applicable.

Competing interests

All authors declare that they have no competing interest.

Author details

¹University Palliative Center, University Hospital Carl Gustav Carus, TU Dresden, Fetscherstraße 74, 01307 Dresden, Germany. ²Department of Psychooncology, National Center for Tumor Diseases (NCT/UCC), Fetscherstraße 74, 01307 Dresden, Germany. ³Core Unit Patient-Reported Outcomes, National Center for Tumor Diseases (NCT/UCC), Fetscherstraße 74, 01307 Dresden, Germany. ⁴Medical Department I, University Hospital Carl Gustav Carus, TU Dresden, Fetscherstraße 74, 01307 Dresden, Germany. ⁵National Center for Tumor Diseases (NCT/UCC), Fetscherstraße 74, 01307 Dresden, Germany.

Received: 14 January 2022 Accepted: 24 April 2022

Published online: 02 May 2022

References

- Ditscheid B, Krause M, Lehmann T, Stichling K, Jansky M, Nauck F, et al. Palliative care at the end of life in Germany: Utilization and regional distribution. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz*. 2020;63(12):1502–10. Available from: <https://europepmc.org/articles/PMC7686196>.
- Prütz F, Saß A-C. Daten zur Palliativversorgung in Deutschland Welche Datenquellen gibt es, und was sagen sie aus? *Bundesgesundheitsblatt - Gesundheitsforsch - Gesundheitsschutz*. 2017;60:26–36.
- Escobar Pinzon LC, Claus M, Zepf KI, Letzel S, Fischbeck S, Weber M. Preference for place of death in Germany. *J Palliat Med*. 2011;14(10):1097–103. Available from: <https://pubmed.ncbi.nlm.nih.gov/22004148/>.
- González-González AI, Schmucker C, Nothacker J, Nury E, Dinh TS, Brueckle M-S, et al. End-of-life care preferences of older patients with multimorbidity: a mixed methods systematic review. *J Clin Med*. 2020;10(1):91. Available from: <https://pubmed.ncbi.nlm.nih.gov/33383951/>.
- Sauer S, Müller R, Rothgang H. Institutionalised dying in Germany: trends in place of death distribution: home, hospitals and nursing homes. *Z Gerontol Geriatr*. 2015;48(2):169–75. Available from: <https://pubmed.ncbi.nlm.nih.gov/24271143/>.
- Gomes B, Calanzani N, Curiale V, Mccrone P, Higginson IJ. Effectiveness and cost-effectiveness of home palliative care services for adults with advanced illness and their caregivers. *Cochrane database Syst Rev*. 2013 Jun 6 [cited 18 Nov 2021];2016(6). Available from: <https://pubmed.ncbi.nlm.nih.gov/23744578/>.
- Peters L, Sellick K. Quality of life of cancer patients receiving inpatient and home-based palliative care. *J Adv Nurs*. 2006;53(5):524–33. Available from: <https://pubmed.ncbi.nlm.nih.gov/16499673/>.
- Higginson IJ, Yi D, Johnston BM, Ryan K, McQuillan R, Selman L, et al. Associations between informal care costs, care quality, carer rewards, burden and subsequent grief: the international, access, rights and empowerment mortality follow-back study of the last 3 months of life (IARE I study). *BMC Med*. 2020 Dec 1 [cited 18 Nov 2021];18(1). Available from: <https://pubmed.ncbi.nlm.nih.gov/33138826/>.
- Gardiner C, Ingleton C, Ryan T, Ward S, Gott M. What cost components are relevant for economic evaluations of palliative care, and what approaches are used to measure these costs? A systematic review. *Palliat Med*. 2017;31(4):323–37. <https://doi.org/10.1177/0269216316670287>.
- Dumont S, Jacobs P, Fassbender K, Anderson D, Turcotte V, Harel F. Costs associated with resource utilization during the palliative phase of care: a Canadian perspective. *Palliat Med*. 2009;23(8):708–17. Available from: <https://pubmed.ncbi.nlm.nih.gov/19837702/>.
- Costa V, Earle CC, Esplen MJ, Fowler R, Goldman R, Grossman D, et al. The determinants of home and nursing home death: a systematic review and meta-analysis. 2016.
- Earle CC, Landrum MB, Souza JM, Neville BA, Weeks JC, Ayanian JZ. Aggressiveness of cancer care near the end of life: is it a quality-of-care issue? *J Clin Oncol*. 2008;26(23):3860–6. Available from: <https://pubmed.ncbi.nlm.nih.gov/18688053/>.
- Alonso-Babarro A, Bruera E, Varela-Cerdeira M, Boya-Cristia MJ, Madero R, Torres-Vigil I, et al. Can this patient be discharged home? factors associated with at-home death among patients with cancer. *J Clin Oncol*. 2011;29(9):1159–67. Available from: <https://pubmed.ncbi.nlm.nih.gov/21343566/>.
- Gamblin V, Prod'homme C, Lecoeuvre A, Bimbai AM, Luu J, Hazard PA, et al. Home hospitalization for palliative cancer care: factors associated with unplanned hospital admissions and death in hospital. *BMC Palliat Care*. 2021 Dec 1 [cited 28 May 2021];20(1). Available from: <https://pubmed.ncbi.nlm.nih.gov/33499835/>.
- Cuppens K, Oyen C, Derweduwén A, Ottevaere A, Sermeus W, Vansteenkiste J. Characteristics and outcome of unplanned hospital admissions in patients with lung cancer: a longitudinal tertiary center study. Towards a strategy to reduce the burden. *Support Care Cancer*. 2016;24(7):2827–35. Available from: <https://pubmed.ncbi.nlm.nih.gov/26816091/>.
- Bouleuc C. La demande d'hospitalisation en urgence des patients atteints de cancer en phase palliative. *Laennec*. 2010;58(2):48–58. Available from: <https://www.cairn.info/revue-laennec-2010-2-page-48.htm?contenu=article>.
- Nilsson J, Blomberg C, Holgersson G, Carlsson T, Bergqvist M, Bergström S. End-of-life care: where do cancer patients want to die? a systematic review. *Asia Pac J Clin Oncol*. 2017;13(6):356–64. Available from: <https://pubmed.ncbi.nlm.nih.gov/28294576/>.
- Hu W, Yasui Y, White J, Winget M. Aggressiveness of end-of-life care for patients with colorectal cancer in Alberta, Canada: 2006–2009. *J Pain Symptom Manage*. 2014;47(2):231–44. Available from: <https://pubmed.ncbi.nlm.nih.gov/23870414/>.
- Costa V, Earle CC, Esplen MJ, Fowler R, Goldman R, Grossman D, et al. The determinants of home and nursing home death: A systematic review and meta-analysis. *BMC Palliat Care*. 2016 Jan 20 [cited 9 Jul 2021];15(1). Available from: <https://pubmed.ncbi.nlm.nih.gov/26791258/>.
- Cohen J, Houttekier D, Onwuteaka-Philipsen B, Miccinesi G, Addington-Hall J, Kaasa S, et al. Which patients with cancer die at home? A study of six European countries using death certificate data. *J Clin Oncol*. 2010;28(13):2267–73. Available from: <https://pubmed.ncbi.nlm.nih.gov/20351336/>.
- Pivodic L, Pardon K, Morin L, Addington-Hall J, Miccinesi G, Cardenas-Turanza M, et al. Place of death in the population dying from diseases indicative of palliative care need: a cross-national population-level study in 14 countries. *J Epidemiol Community Health*. 2016;70(1):17–24. Available from: <https://pubmed.ncbi.nlm.nih.gov/26202254/>.
- Woitha K, Garralda E, Martin-Moreno JM, Clark D, Centeno C. Ranking of palliative care development in the countries of the European union. *J Pain Symptom Manage*. 2016;52(3):370–7. Available from: <https://pubmed.ncbi.nlm.nih.gov/27287622/>.

23. Being a Caregiver | Johns Hopkins Medicine. [cited 20 Dec 2021]. Available from: <https://www.hopkinsmedicine.org/health/caregiving/being-a-caregiver>
24. Wetzstein M, Rommel A, Lange C. Pflegende Angehörige–Deutschlands größter Pflegedienst. 2015 [cited 19 Nov 2021]; Available from: <https://edoc.rki.de/handle/176904/3137>
25. Toukhsati S, Hare D. Towards optimal heart failure care: couples-oriented strategies to improve patient adherence and health outcomes. *Curr Cardiol Rev.* 2016;12(3):243–8. Available from: <https://pubmed.ncbi.nlm.nih.gov/27280305/>.
26. Pinquart M, Sörensen S. Differences between caregivers and noncaregivers in psychological health and physical health: a meta-analysis. *Psychol Aging.* 2003;18(2):250–67. Available from: <https://pubmed.ncbi.nlm.nih.gov/12825775/>.
27. O'Toole MS, Zachariae R, Renna ME, Mennin DS, Applebaum A. Cognitive behavioral therapies for informal caregivers of patients with cancer and cancer survivors: a systematic review and meta-analysis. *Psychooncology.* 2017;26(4):428–37. <https://doi.org/10.1002/pon.4144>.
28. Longacre ML, Applebaum AJ, Buzaglo JS, Miller MF, Golant M, Rowland JH, et al. Reducing informal caregiver burden in cancer: evidence-based programs in practice. *Transl Behav Med.* 2018;8(2):145. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6257028/>.
29. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)—A metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform.* 2009;42(2):377–81. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/18929686>.
30. Harris PA, Taylor R, Minor BL, Elliott V, Fernandez M, O'Neal L, et al. The REDCap consortium: building an international community of software platform partners. *J Biomed Inform.* 2019;95:103208. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/31078660>.
31. The WHOQOL Group. Development of the world health organization WHOQOL-BREF quality of life assessment. *Psychol Med.* 1998;28(3):551–8. Available from: <https://pubmed.ncbi.nlm.nih.gov/9626712/>.
32. Mehnert A, Müller D, Lehmann C, Koch U. Die deutsche version des NCCN Distress-Thermometers: Empirische prüfung eines screening-instruments zur erfassung psychosozialer belastung bei krebspatienten. *Zeitschrift für Psychiatrie Psychol und Psychother.* 2006;54(3):213–23.
33. McGeorge P, Holmes D, Bernardi S, Vincent S, Health Services H, Centre for Urban Health B, et al. Recognising and responding to deterioration in mental state A scoping review. 2014.
34. Spitzer RL, Kroenke K, Williams JBW, Löwe B. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med.* 2006;166(10):1092–7.
35. Kroenke K, Spitzer RL, Williams JBW. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med.* 2001;16(9):606. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1495268/>.
36. Downar J, Goldman R, Pinto R, Englesakis M, Adhikari NKJ. The “surprise question” for predicting death in seriously ill patients: a systematic review and meta-analysis. *CMAJ.* 2017;189(13):E484–93.
37. First MB, Williams JBW, Karg RS, Spitzer RL. User's Guide to Structured Clinical Interview for Dsm-5 Disorders (Scid-5-cv): Clinician Version Title : User's Guide to Structured Clinical. 2015 [cited 24 Nov 2021 Nov 24];5–7. Available from: <https://jrwwq1tbn01.storage.googleapis.com/MTU4NTYyNTI0OA==01.pdf>
38. Philipp R, Kalender A, Härter M, Bokemeyer C, Oechsle K, Koch U, et al. Existential distress in patients with advanced cancer and their caregivers: Study protocol of a longitudinal cohort study. *BMJ Open.* 2021 Apr 24 [cited 2021 June 4];11(4). Available from: <https://pubmed.ncbi.nlm.nih.gov/33895716/>

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

